INDEX

| Alabama plants104 | Botonical Gleanings263 |
|--------------------------------|---------------------------------|
| Allionia aggregata269 | Bromus inermis265 |
| Gladewitzii 268 | |
| Nyctaginia 268 | Capnorchis 62 |
| Alnus crispa267 | Carex aristata266 |
| mollis268 | echinata266 |
| Alsine rubra 269 | exilis266 |
| Amblema phiata135 | gynocrates266 |
| Amygdalus persica272 | hirta266 |
| Anethum graveoleus275 | |
| Anodonta cataracta281 | Castilleja coccinea276 |
| implicata281 | Cerastium longepedunculatum 269 |
| grandis footiana138 | Conchology |
| Anodontoides f. s140 | Collection of shells 6 |
| Archeology 49 | Conocephalum conicum222 |
| Arctium Lappa279 | Cosmos bipinnatus278 |
| Arethusa bulbosa269 | Cryptophila pudica108, 117 |
| Arisaema triphyllum40, 46, 266 | Cunila 61 |
| Asclepias Sullivantii275 | Cururu 63 |
| verticillata275 | |
| Aster ericoides277 | Darwinism211 |
| leavis277 | Death rate in animals153 |
| Atriplex argentea268 | Deschampsia caespitosa265 |
| Audubon societies 65 | Dilepyrum, type of 33 |
| | Dimorphism, seasonal 41 |
| Biology, animal 39 | Diplotaxis tenuifolia271 |
| Birds- | Disporum Cahnae267 |
| of Bankson Lake260 | Dogfish159 |
| geography of 65 | Drosena linearis272 |
| of Floyd County189, 231 | |
| migration74, 81 | Echinoid, Devonian219 |
| of Notre Dame257, 283 | Eleocharos rostellata265 |
| of Northfield 73 | Elliptio dilatus skerkii137 |
| of River Forest244 | Eragrostis Eragrostis265 |
| of Washington 35 | Erigeron Canadense277 |
| of Watertown172 | Eriophorium Virg265 |
| Blephariglottis ciliaris267 | Erucastrum Gallicum271 |
| Borbonia 63 | Euglena viridis149 |

INDEX

| Euphorbia corollata | 273 | Lasmigonia costata erigar | nensis 138 |
|------------------------------|-----|---------------------------|------------|
| dentata | 273 | Leptodea f | 139 |
| | | Lespedeza capitata | 272 |
| Falcata Pitcheri | | Ligumia | 140 |
| Fauna, mussel | | Lilium philadelphicum | 266 |
| Fish, death rate | | | |
| Flower bed, wild | 164 | Marchantiaceae | |
| Floyd County | | Mentha Cardiaca | |
| Forests | | Migration records | |
| Industries | 192 | Moehringia lateriflora | 269 |
| Population | 192 | Monarda fistulosa | |
| Roads | | Monotropoid, new genus . | 104 |
| Topography | 190 | Monotropsis | 113 |
| Forma alba | 276 | Muhlenbergia racemosa | |
| lutescens | 276 | Mussel, collection of | 7 |
| Fraxinus Michanxii | 275 | fauna, type of | 131 |
| Fusconaja flava paurola | 134 | fisheries | |
| | | gills | 89 |
| Gambusia holbrooki | 160 | Lake Pepin | 16 |
| Gardenia | 34 | | |
| Gentiana | 45 | Naiades | 89, 281 |
| Geum canadense | 272 | Naias flexilis | |
| virginianum | | Newton, Alfred | 69 |
| Gills, | | Nigella Damascena | |
| histology of | 90 | Nomenclature, botanical | 33, 61 |
| method of study | | Nortonechinus primus | |
| Gniabara | | Nymphozanthus advena | 270 |
| | | variegatus | 270 |
| Helianthus scaberrimus | 278 | variegavas | |
| lenticularis | | Oenothera biennis | 274 |
| Heliopsis helanthoides | | cruciata | 273 |
| scabra | | parviflora | |
| Hieracium florentinum | | Osmerus eperalanus | 159 |
| Histology of gills | | | |
| Hippoglossoides platessoides | | Panicum barbulatum | |
| Humulus japonicus | | Werneri | 265 |
| Lupulus | | Paracelsus | 39 |
| Dupulus | | Persea | 63 |
| Implements, evolution of | 40 | Physalis heterophyla | 276 |
| Isanthus brachiatus | | Plaice, death rate | 157 |
| Isopyrum beternatum | | Plantago lanceolata | 276 |
| Loopyrum beternatum | 411 | Pleurobema coccineum | 136 |
| Juniperius virginiana | 969 | Pleuronectes cynoglossus | |
| oumperius virginiana | 203 | platessa | 157 |
| Lampsilis | 142 | Poa sylvestris | 265 |
| Lampsilis | | Polygala ambigua | 273 |
| | | | |

HHI

INDEX

7 ---131 --- 2 --- 89 --- 16

9, 281 ___265

__274 __278 __278 __159 __265 __39 __63 __276 __157 __276 __136 __157 __157 __265 __273

| Potamogeton alpinus | 269 | Salmo salar | 159 |
|----------------------|-----|------------------------|-----|
| americanus | | Sanguisorba Canadensis | 272 |
| amphifolius | 264 | Sassafras Sassafras | 271 |
| augustifolius | 264 | Scutellaria parvula | 276 |
| foliosus | 265 | Shells, collection of | 6 |
| gramineus | 264 | rate of growth | 129 |
| lucens | | Silene latifolia | 270 |
| perfoliatus | 264 | Silphium lacinatum | 278 |
| Poterium Sanguisorba | | Sinapis alba | 271 |
| Preissia commutata | 223 | Smelt | 159 |
| Primula Mistassinica | 275 | Solidago graminifolia | 277 |
| Proptera | 139 | Sonchus arvensis | 280 |
| Prunus Avium | | Spinax niger | |
| Pussila, A. T. | 42 | - | |
| Radicula sylvestris | 271 | Tools, evolution of | 49 |
| Ranunculus acris | | Top minnow | 160 |
| Robin | | Tragopogen major | 279 |
| Rudbeckia laciniata | | parrifolius | 279 |
| triloba | | pratensis | 279 |
| Rummex patientia | | Tusilago Farfara | 278 |
| Sagina procumbens | 269 | Vernonia interior | 277 |
| Salix alba | 267 | Vicia sparsiflora | 272 |
| condida | 267 | Viola palens | 273 |
| pedicellaris | 267 | rostrata | 273 |

χι